

Half-lives of Double β^+ -decay with Two Neutrinos

Yuejiao Ren¹ and Zhongzhou Ren^{*1,2}

¹*Department of Physics, Nanjing University, Nanjing 210093, China*

²*Center of Theoretical Nuclear Physics,*

National Laboratory of Heavy-Ion Accelerator, Lanzhou 730000, China

Abstract

Nuclear double β^- -decays with two neutrinos were observed for many years and a systematic law describing the relation between their half-lives and decay energies was also proposed recently [Phys. Rev. C89, 064603 (2014)]. However, double β^+ -decay ($\beta^+\beta^+$) with emission of both two positrons and two neutrinos has not been observed up to date. In this article, we perform a systematic analysis on the candidates of double β^+ -decay, based on the 2012 nuclear mass table. Eight nuclei are found to be the good candidates for double β^+ -decay and their half-lives are predicted according to the generalization of the systematic law to double β^+ -decay. As far as we know, there is no theoretical result on double β^+ -decay of nucleus ^{154}Dy and our result is the first prediction on this nucleus. This is also the first complete research on eight double β^+ -decay candidates based on the available data of nuclear masses. It is expected that the calculated half-lives of double β^+ -decay in this article will be useful for future experimental search of double β^+ -decay.

Key words : double β^+ -decay with two neutrinos, half-lives, systematic law, weak interaction.

PACS numbers: 23.40.-s, 23.40.Bw

* Corresponding author, zren@nju.edu.cn

I. INTRODUCTION

Nuclear double β -decay with two neutrinos is a rare and exotic process occurring in nuclei with long lifetimes. Since Goeppert-Mayer predicted that there is double β -decays in 1935 [1], many researches have been carried out on double β -decay with two neutrinos or without neutrinos [2–10]. Plenty of results can be found in the references of the articles [11–23]. Although much experimental effort has been devoted to double β -decay, only the half-lives of eleven nuclei with double β^- -decays were obtained and the half-life of a single nucleus with double electron capture (ECEC) was measured due to the difficulty of experimental detection in this field [8]. On double β^+ decay ($\beta^+\beta^+$), experimental data of definite half-lives are still not available [8]. Theoretically there are many successful calculations on the double β^- -decay half-lives [2–4, 12–23]. However, calculations on the double β^+ -decay half-lives are much less as they are compared with those on double β^- -decay [2–4, 12–23]. Haxton et al. estimated the half-lives of six nuclei from mass number $A=78$ (^{78}Kr) to mass number $A=136$ (^{136}Ce) and showed that their half-lives for double β^+ -decay ranged approximately from 10^{26} years to 10^{31} years [2]. Since the calculations from Haxton et al. [2], thirty years have passed and the data of nuclear masses have been updated significantly [2, 9, 10]. Therefore, it is necessary to calculate the half-lives of double β^+ -decay with new nuclear mass tables because the half-lives are very sensitive to the decay energies. Recently, Suhonen made the theoretical investigation of double β^+ -decay for two nuclei, ^{78}Kr [19] and ^{96}Ru [20]. As definite double β^+ -decay half-lives are not available in experiments now, it is interesting to make a complete calculation on the possible candidates of double β^+ -decay and to predict new candidates for future experiments.

II. NUMERICAL RESULTS AND DISCUSSIONS ON THE DOUBLE β^+ -DECAY HALF-LIVES

Based on the 2012 nuclear mass table [9, 10], we calculate the decay energy of all possible double β^+ -decay emitters and pay special attention to the parent nuclei where they are denoted with a symbol ($2\beta^+?$) in the table. The number of possible double β^+ -decay emitters is 40 in the mass table [9, 10] and their mass number ranges from $A=36$ (^{36}Ar) to $A=252$ (^{252}Fm) [9, 10]. In accordance with textbooks and published articles, the decay

energy of double β^+ -decay is defined as

$$Q_{2\beta}(\text{MeV}) = [M(A, Z) - M(A, Z - 2) - 4 M_e] C^2 \quad (1)$$

where $M(A, Z)$ is the mass excess of the parent nucleus and $M(A, Z-2)$ is the mass excess of the daughter nucleus. M_e is the rest mass of an electron (or a positron) and C is the speed of light in vacuum ($C = 1$ in natural units). Our numerical calculations for the 40 possible emitters shows that many of them have negative decay energy to double β^+ -decay or have approximately zero decay energy. Only eight of them have significantly positive energies for double β^+ -decay. We list the eight emitters (parent nuclei) and their daughter nuclei in Table 1, together with the decay energy and the isotopic abundance (IS) of parent nuclei (or their α -decay half-lives (T^α) when α -decay is observed).

TABLE I: The decay energy of double β^+ -decay candidates from the 2012 nuclear mass table where the decay energy is defined as $Q_{2\beta} = [M(A, Z) - M(A, Z - 2) - 4 M_e] C^2$ (MeV). We also list the isotopic abundance (IS) of parent nuclei or their α -decay half-lives (T^α) when α -decay is observed.

The experimental data of nuclear masses and isotope abundance are from references [9, 10].

Parent	Daughter	M(A,Z) (MeV)	M(A,Z-2) (MeV)	$Q_{2\beta}$ (MeV)	IS or T^α
^{78}Kr	^{78}Se	-74.180	-77.026	0.802	IS=0.355%
^{96}Ru	^{96}Mo	-86.079	-88.794	0.671	IS=5.54%
^{106}Cd	^{106}Pd	-87.132	-89.907	0.731	IS=1.25%
^{124}Xe	^{124}Te	-87.661	-90.525	0.820	IS=0.095%
^{130}Ba	^{130}Xe	-87.262	-89.880	0.574	IS=0.106%
^{136}Ce	^{136}Ba	-86.509	-88.887	0.334	IS=0.185%
^{148}Gd	^{148}Sm	-76.269	-79.336	1.023	$T^\alpha=70.9$ y
^{154}Dy	^{154}Gd	-70.394	-73.705	1.267	$T^\alpha=3.0$ My

Table 1 shows that all eight parent nuclei have positive decay energies for double β^+ -decay. From ^{78}Kr to ^{136}Ce , they are naturally occurring isotopes and their decay energy is lower than 1 MeV. For nuclei ^{148}Gd and ^{154}Dy , they are unstable for α -decay but their α -decay half-lives are long ($T^\alpha=70.9$ years (y) or $T^\alpha=3.0$ My) [9, 10]. Especially their decay energies for double β^+ -decay are higher than 1 MeV. So the last two nuclei (^{148}Gd and ^{154}Dy) could be very interesting to observe double β^+ -decay because decay half-lives are

very sensitive to decay energies.

For the purpose of calculating the double β^+ -decay half-lives of the above eight nuclei, we will generalize the systematic law of double β^- -decay half-lives [24] to the case of double β^+ -decay. It is proposed [24] that there is a new systematic law between the half-lives and decay energies for double β^- -decay with two neutrinos

$$\lg T_{1/2}(Ey) = (a - 2 \lg(2\pi Z/137) + S) / Q_{2\beta}(MeV) \quad (2)$$

where the constant a is obtained by fitting the experimental data of double β^- -decay and its value is $a = 5.843$ [24]. The physical meaning of a is related to the square of the strength of the weak interaction which leads to the instability of a nucleus. Z is the charge number of the parent nucleus and the second term in the right side of Eq. (2) is the effect of the Coulomb field on double β^- -decay half-lives due to the emission of two electrons [24]. $S = 2$ when the neutron number of parent nuclei is a magic number and $S = 0$ when the neutron number of parent nuclei is not a magic number [24].

In order to generalize the law of Eq.(2) from double β^- -decay to that of double β^+ -decay, we analyze the contribution of the Coulomb field on half-lives. It is well known from textbooks [25–27] and from previous researches [28–33] that the effect of the Coulomb field in single β^+ -decay is opposite as compared with that in single β^- -decay. The effect of the Coulomb field will shorten the half-lives of a β^- -decay but it will prolong the half-lives of a β^+ -decay when the decay energy is fixed. This is due to the opposite charge between a positron and an electron. These are also confirmed by numerical calculations of β -decay half-lives [28–33]. It is natural to extend this idea from single β -decay to double β -decay. Therefore, the systematic law of double β^+ -decay half-lives is

$$\lg T_{1/2}(Ey) = (a + 2 \lg(2\pi Z/137)) / Q_{2\beta}(MeV) \quad (3)$$

Here the parameter a of eq.(3) is the same value as that in Eq.(2). The second term of eq.(3) is the same value as that in Eq.(2) but with opposite contribution due to different signs of charges between positron and electron. In Eq. (2) the last term S is from the contribution of neutron closed shell when the neutron number of parent nucleus is a magic number for double β^- -decay. However, for double β^+ -decay, it is difficult to occur due to the requirement of decay energy and there are only eight suitable candidate nuclei in Table 1. The proton number of the eight nuclei in Table 1 is not a magic number and therefore

it is natural to choose $S = 0$. Finally we obtain the analytical formula Eq.(3) to calculate the half-lives of double β^+ -decay, without introducing new adjusting parameters. So it can be concluded that the number of adjusting parameters in this article is the minimum, with the same spirit as that in our previous research [24].

We use Eq.(3) to calculate the half-lives of double β^+ -decay and the numerical results are drawn in Figure 1 and listed in Table 2. In Fig.1, the X-axis is the decay energy (MeV) and the Y-axis is the logarithm of the double β^+ -decay half-lives (Ey) where $Ey = 10^{18}$ years. Fig.1 shows that the calculated half-lives are very sensitive to the variation of the decay energies. Although decay energies vary in a narrow range from 0.334 MeV (^{136}Ce) to 1.267 MeV (^{154}Dy), the change to the half-lives is approximately a factor of 10^{15} . This strongly suggests that double β^+ -decay with a higher decay energy is better for observation. We recommend nuclei ^{154}Dy and ^{148}Gd as good candidates to observe double β^+ -decay.

Now we make a detailed discussion on the calculated results of Table 2. In Table 2, Column 1 represents the parent nuclei and column 2 represents experimental decay energy. Column 3 denotes the logarithm of the calculated double β^+ -decay half-lives with equation (3). Column 4 is our calculated double β^+ -decay half-lives, which is convenient for comparison with future experiments. The fifth column provides the calculated results from Suhonen [19, 20] and they are also listed for comparison. It is seen from the comparison of our results (column 4) and other results (the last column) that the two sets of theoretical half-lives are close for nuclei ^{78}Kr and ^{96}Ru although different calculation approaches were used. There was an old calculation [12] in 1991 on seven nuclei but it is not suitable for comparison with our results because the experimental data of nuclear masses have been updated twice. This is related to different inputs of decay energies between the old calculation and that of this manuscript. So reasonable agreement is reached for different calculations. This is good for future experiments. It is widely accepted that different approaches are useful for the further development of the field. From the last two rows of Table 2, it is seen that the half-lives of ^{148}Gd and ^{154}Dy are shorter due to their higher double β^+ -decay energies. Because both ^{148}Gd and ^{154}Dy have been observed to have α -decay and their α -decay half-lives are long ($T^\alpha=70.9$ years for ^{148}Gd and $T^\alpha=3.0$ My for ^{154}Dy) [9, 10, 34, 35], it is interesting to detect the double β^+ -decay from the two nuclei. We can estimate a branch ratio [9, 10, 34] from the half-lives of double β^+ -decay (Table 2) and α -decay (Table 1). The branching ratio (BR) between double β^+ -decay and α -decay is

$BR = T^\alpha/T_{1/2}(theor) = (3 \times 10^6) / (2.35 \times 10^5 \times 10^{18}) = 1.28 \times 10^{-17}$ for ^{154}Dy . This value can be used for reference of future experiments to search the double β^+ -decay of ^{154}Dy .

TABLE II: The double β^+ -decay half-lives of even-even isotopes calculated with new systematic law (T_{theor}) and the corresponding logarithms (lgT_{theor}). The units of the half-lives are Ey (10^{18} years). The experimental decay energies of nuclei ($Q_{2\beta}$ (MeV)) are also listed in the table. The calculated half-lives from another group [19, 20] are listed in the last column for comparison.

Nuclei	$Q_{2\beta}(\text{MeV})$	$lgT_{1/2}(theor)$	$T_{1/2}(theor)$ (Ey)	$T_{1/2}(other)$ (Ey)
^{78}Kr	0.802	7.828	6.73×10^7	$(4.94 - 15.8) \times 10^7$ [19]
^{96}Ru	0.671	9.616	4.13×10^9	$(1.2 - 10) \times 10^8$ [20]
^{106}Cd	0.731	8.930	8.51×10^8	
^{124}Xe	0.820	8.086	1.22×10^9	
^{130}Ba	0.574	11.606	4.04×10^{11}	
^{136}Ce	0.334	20.037	1.09×10^{20}	
^{148}Gd	1.023	6.625	4.22×10^6	
^{154}Dy	1.267	5.371	2.35×10^5	

III. SUMMARY

In summary, we collect the experimental data of possible double β^+ -decay nuclei based on the 2012 nuclear mass table and find that eight nuclei have significantly positive decay energies. A systematic calculation of double β^+ -decay half-lives for the eight nuclei is carried out with the analytical formula being a natural generalization of the systematic law from double β^- -decay to double β^+ -decay. Numerical results show that the half-lives of double β^+ -decay of nuclei ^{148}Gd and ^{154}Dy are shorter than other nuclei and they can be very interesting for future experimental observation of double β^+ -decay. The branching ratio (BR) between double β^+ -decay and α -decay is also estimated for ^{154}Dy . Our result on nucleus ^{154}Dy is the first prediction as far as we know. This is a complete calculation on eight double β^+ -decay nuclei, which will be useful for future experimental researches [36].




FIG. 1: (Color online) Logarithms of the theoretical double β^+ -decay half-lives for ground-state transitions of eight even-even parent nuclei from ^{78}Kr to ^{154}Dy . The units of the half-lives are Ey (10^{18} years). X-axis is the decay energy (MeV). The half-lives of double β^+ -decay of nuclei ^{148}Gd and ^{154}Dy are shorter than other nuclei and they could be very interesting for future experimental observation of double β^+ -decay.

IV. ACKNOWLEDGEMENTS

This work is supported by National Natural Science Foundation of China (Grant Nos. 11035001, 10975072, 11120101005, 11175085, 11375086), by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

-
- [1] M. Goeppert-Mayer, Phys. Rev. **48**, 512 (1935).
 - [2] W. C. Haxton, G. J. Stephenson, Prog. Part. Nucl. Phys. **12**, 409 (1984).
 - [3] H. Ejiri, Phys. Rep. **338**, 265 (2000).
 - [4] N. Ackerman, B. Aharmim, M. Auger, D. J. Auty, P.S. Barbeau et al., Phys. Rev. Lett. **107**, 212501 (2011).

- [5] A. Gando, Y. Gando, H. Hanakago et al., Phys. Rev. C **85**, 045504 (2012) .
- [6] R. Arnold, C. Augier, J. Baker, A. S. Barbash, A. Basharina-Freshville, S. Blondel et al., Phys. Rev. Lett. **107**, 062504 (2011).
- [7] A. S. Barabash, Phys. Rev. C **81**, 035501 (2010).
- [8] R. Saakyan, Annu. Rev. Nucl. Part. Sci. **63**, 503 (2013).
- [9] G. Audi,, F. G. Kondev, M. Wang, B. Pfeiffer, X. Sun, J. Blachot, M. MacCormick, Chin. Phys. C **36**, 1157 (2012).
- [10] M. Wang, G. Audi, A. H. Wapstra, F. G. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, Chin. Phys. C **36**, 1603 (2012).
- [11] S. R. Elliott, A. A. Hahn, M. K. Moe, Phys. Rev. Lett. **59**, 2020 (1987).
- [12] A. Staudt, K. Muto, H. V. Klapdor-Kleingrothaus, Phys. Lett. B **168**, 312 (1991).
- [13] A. A. Raduta, C. M. Raduta, Phys. Lett. B **647**, 265 (2007).
- [14] F. T. Avignone III, S. R. Elliott, J. Engel, Rev. Mod. Phys. **80**, 481 (2008).
- [15] M. Agostini et al., J. Phys. G **40**, 035110 (2013).
- [16] F. Simkovic, V. Rodin, A. Faessler, P. Vogel, Phys. Rev. C **87**, 045501 (2013).
- [17] A. Faessler and F. Simkovic, J. Phys. G **24**, 2139 (1998).
- [18] E. Caurier, F. Nowacki, A. Poves, Phys. Lett. B **711**, 62 (2012).
- [19] J. Suhonen, Phys. Rev. C **87**, 034318 (2013).
- [20] J. Suhonen, Phys. Rev. C **86**, 024301 (2012).
- [21] J. Suhonen, O. Civitarese, Phys. Rep. **300**, 123 (1998).
- [22] P. Vogel and M. R. Zirnbauer, Phys. Rev. Lett. **57**, 3148 (1986).
- [23] J. D. Vergados, Phys. Rep. **361**, 1 (2002).
- [24] Yuejiao Ren and Zhongzhou Ren, Phys. Rev. C **89**, 064603 (2014) .
- [25] M. A. Preston, Physics of the Nucleus, (Addison-Wesley, Reading, Massachusetts,1962).
- [26] A. De Shalit and H. Feshbach, Theoretical Nuclear Physics, Volume 1,(John Wiley and Sons, Newy York, 1974).
- [27] J. D. Walecka, Theoretical Nuclear and Subnuclear Physics, Imperial College Press (London) and World Scientific (Singapore), 2004.
- [28] Dongdong Ni and Zhongzhou Ren, J. Phys. G **39**, 125105 (2012).
- [29] Dongdong Ni and Zhongzhou Ren, J. Phys. G **41**, 025107 (2014).
- [30] Hantao Li and Zhongzhou Ren, J. Phys. G **40**, 105110 (2013).

- [31] Hantao Li and Zhongzhou Ren, Sci China-Phys Mech Astron **57**, 1005 (2014) .
- [32] Xiaoping Zhang and Zhongzhou Ren, Phys. Rev. C **73**, 014305 (2006) .
- [33] P. Moeller, J. R. Nix, and K. Kratz, At. Data Nucl. Data Tables **66**, 131 (1997) .
- [34] Yuejiao Ren and Zhongzhou Ren, Phys. Rev. C **85**, 044608 (2012) .
- [35] Yibin Qian and Zhongzhou Ren, Sci China-Phys Mech Astron **56**, 1520 (2013) .
- [36] Xiguang Cao et al., Sci China-Phys Mech Astron **57**, 1476 (2014) .